



Does the lower extremity alignment affect the risk of falling?

Gülnur Taşçı Bozbaş , Gülcan Gürer 

Department of Physical Medicine and Rehabilitation, Medical Faculty of Adnan Menderes University, Aydın, Turkey

Received: Mart 06, 2017 Accepted: September 26, 2017 Published online: October 03, 2017

ABSTRACT

Objectives: This study aims to investigate the effects of knee and foot alignments on the risk of falling.

Patients and methods: Between April 2016 and December 2016, a total of 74 individuals (24 males, 50 females; mean age 32.2±4.9 years; range 18 to 65 years) were included in the study. The knee Q angle and Chippaux-Smirak Index (CSI), Arch Index, and foot progression angle (FPA) evaluated by pedobarography were used for the assessment of the lower extremity alignment. The fall risk was evaluated by the Fall Index, Fourier 56 Index (F56), and Stability Index.

Results: The fall index was found to be correlated with the Q angle, CSI, the Arch index, and FPA ($p<0.05$). Q angle, Arch Index, and FPA which were explained 40% of the variance of the fall index. The Q angle was correlated with F56 and the stability index at the most position ($p<0.05$). The CSI was correlated with the F56 and the stability index at two and three positions, respectively ($p<0.05$); however, the Arch Index and FPA were not correlated with the F56 and Stability Index at any of the eight positions ($p>0.05$). According to the categorical regression analysis, the Q angle was the most effective on the F56 and Stability Index.

Conclusion: Our study results suggest that lower extremity malalignment increases the risk of falling. We believe that the risk of falling can be decreased by the reduction of these malalignments and, thus, mortality and morbidity associated with the fall can be reduced as well.

Keywords: Arch Index; Chippaux-Smirak Index; falling; foot progression angle; pedobarography; Q angle.

Postural control is the spatial control of the body position with the aim of providing balance.^[1] This control is maintained by regulating the peripheral stimuli from the somatosensory, visual, and vestibular system in the central nervous system and producing muscular motor responses. As a result, falling is prevented by balancing the movement of the center of gravity on the plantar surface at the weight-bearing position.^[2] Center-of-pressure (COP) excursions are calculated through the ground reaction force and are the indicative of postural control.^[3] Stability disorders may cause disabilities and even falls which can lead to serious morbidities.^[4-6] Falling is one of the most critical events preventing the participation of the individual from social life and reducing the capability of self-care.^[7] Therefore, it is an important public health issue regarding its medical and economic consequences. There are at least two risk factors for falling and the risk increases with increasing risk

factors in individuals with the history of falling.^[8] Therefore, identifying the variables affecting postural stability is of utmost importance to prevent falls.

The lower extremity plays an important role in various static and dynamic activities such as standing, walking, running, and swimming.^[9] Therefore, the alignment disorders of this region can affect stability. One of the commonly used parameters in the assessment of biomechanical alignment of the lower extremity is the Q angle of the knee. This angle was first defined by Brattström.^[10] Using this approach, 10° to 20° for females and 6° to 10° for males are accepted normal. The deviations of the Q angle are named as genu varum or genu valgum of the knee joint. The Chippaux-Smirak Index (CSI), Arch Index, and the Foot Progression Angle (FPA) are useful for the biomechanical evaluation of the alignment of the foot.^[11] The changes in the CSI and Arch Index indicate flat foot or pronation and cavus foot or supination.

Corresponding author: Gülnur Taşçı Bozbaş, MD. Adnan Menderes Üniversitesi Tıp Fakültesi, Fiziksel Tıp ve Rehabilitasyon Anabilim Dalı, 09010 Aydın, Turkey.
e-mail: gulnurtb@yahoo.com

Cite this article as:

Taşçı Bozbaş G, Gürer G. Does the lower extremity alignment affect the risk of falling? Turk J Phys Med Rehab 2018;64(2):140-147.

It was presented at 4th Physical Medicine and Rehabilitation Orthopedic and Traumatology days, December 15-17, 2016, Izmir, Turkey

In addition, FPA is the angle between the foot and the progression line, and values between -3° and $+20^\circ$ are accepted normal.^[12] Higher values indicate “out-toeing,” while lower values indicate “in-toeing”.^[13]

Several studies have shown that change in the alignment of the lower extremity plays a role in the injury formation.^[14-17] In addition, the effects of some changes in the alignment of the foot on the postural excursions have been also investigated.^[3,18-20] However, the direct effect of lower extremity alignment on the risk of falling has not been emphasized. Indeed, the knee and foot alignments play a significant role in the lower extremity functions and daily living activities. Therefore, in the present study, we aimed to investigate the effects of foot and knee alignments on the risk of falling in healthy individuals.

PATIENTS AND METHODS

In this cross-sectional study, a total of 217 individuals between the ages of 18 and 65 years were screened between April 2016 and December 2016. Those who had any vestibular and visual problems and neurological or orthopedic disorders in the lower extremity, who underwent ankle, foot, and knee surgery, those with lower extremity pain leading to walking difficulty, and those who received medications disrupting the balance control were excluded. As a result, a total of 74 individuals (24 males, 50 females; mean age 32.2 ± 4.9 years; range 18 to 65 years) were included in the study. The study was approved by the institutional Ethics Committee (2014/499). A written informed consent was obtained from each participant. The study was conducted in accordance with the principles of the Declaration of Helsinki.

Demographic characteristics such as age, sex, height, and weight and the history of falling of all participants were recorded. The body mass index (BMI) was calculated as weight in kilograms divided by the square of height in meters (kg/m^2). Postural sway and risk of falling were assessed using the tetra-ataxiometric posturography (Tetrax*-Sunlight medical Ltd., Ramat Gan, Israel). The device was calibrated before the study. The participants were assessed in eight different positions by spending 32 sec on each position (Table 1). The Fall Index, Fourier Index, and Stability Index scores were calculated from the obtained data by Tetrax* software (Tetrax-Sunlight medical Ltd., Ramat Gan, Israel). The Fall Index is calculated with a proprietary algorithm, which provides varying weights to different parameters for the balance results. It is an indicator of the risk of

falling and divided into three groups: 0-36 indicate minimal risk of falling; 37-58 indicate moderate risk; and 59-100 indicate high risk.^[21] The Fourier Index is an analysis of postural sway using different frequencies. The Tetrax* software subdivides postural sways into four categories as follows: F1: low frequencies (below 0.1 Hz), F2-4: lower-to-medium frequencies (0.1-0.5 Hz), F5-F6: high-to-medium frequencies (0.5-1 Hz), and F7-F8: high frequencies (above 1 Hz). The sway at a high-to-medium range (0.5-1Hz) (F5, F6) indicates somatosensory response mediated by the motor function of the lower extremities. The Stability Index measures the amount of sway over the four plates and is an indicator of the general stability. Higher index scores reflect more unstable posture.^[21]

The Q angle was used to evaluate the biomechanical alignment of the knee. It was measured with a lengthened arm 360° plastic universal goniometer. The participant was positioned in supine position with toes pointing vertically (knee extended and quadriceps muscle relaxed). The center of the patella, the tibial tuberosity, and the anterior superior iliac spine were labeled with a marker. The pivot of the goniometer was placed on the center of the patella. The long arms of the goniometry were placed on the tibial tuberosity and the anterior superior iliac spine. The intersected angle was regarded as the Q angle.^[22] The measurement was performed three times in the dominant side of the knee and mean values were calculated.

The biomechanical properties of the foot were measured using pedobarography (RSscan International, Olen, Belgium). The participant was asked to stand with bare feet on a 0.5 m pressure-sensitive platform and to walk with normal steps across this platform embedded in the middle of a 3-m-long walkway.^[23] The measurements were repeated twice and mean value of the two measurements was recorded as the final score and included in the analysis. In dynamic measurements, the foot axis was, then, divided into three equal lengths to divide the footprint into a heel, midfoot, and forefoot area. The Arch Index was calculated as the ratio midfoot area divided by the whole footprint area, excluding toes. This index is a useful indicator of foot type to be a valid clinical means.^[24] The CSI is measured by dividing the value of the narrower zone of the midfoot by the value of the parallel line on the wider zone of the forefoot and multiplying by 100.^[11] In the static measurement, the foot axis line from the second toe to the middle of the heel was drawn in the footprint. The FPA was derived as the angle in the transverse plane between the foot vectors (second metatarsal to the posterior middle calcaneous).^[12]

Table 1. Eight positions on the tetra-ataxiometric posturography

Statistical analysis

Statistical analysis was performed using the IBM SPSS version 22.0 software (IBM Corp., Armonk, NY, USA). Descriptive statistics were expressed in mean and standard deviation (SD), and median (min-max) values. Compliance with the normal distribution of the quantitative data was examined using the Shapiro-Wilk test. The correlations between the quantitative variables were analyzed using the Pearson's correlation

Table 2. Descriptive statistics for Q angle, Chippaux-Smirak Index, Arch Index, and foot position angle

	Mean±SD	Median	Range
Fall index	38.5±28.5	32.0	6-100
Q angle	9.6±3.2	8.5	4-17
Chippaux-Smirak index	30.7±8.7	31.0	15-53
Arch index	0.3±0.1	0.32	0.21-0.42
Foot position angle	12.9±7.1	12.6	0.09-33.2

SD: Standard deviation.

analysis. The categorical regression analysis was used to identify the factors which would affect the Fall Index, F56, and Stability Index. A *p* value of <0.05 was considered statistically significant.

RESULTS

Of the participants, the mean BMI was 26.1±4.3 (median: 25.8) kg/m². Only seven participants had a history of falling. The mean and median values of the Fall Index, Q angles, CSI, Arch Index, and FPA are shown in Table 2. The correlation between age, sex, and BMI and the Fall Index, Q angle, CSI, Arch Index, and FPA was analyzed. The age and BMI were not correlated with the Fall Index, Q angle, CSI, Arch Index, and FPA (*p*>0.05). Furthermore, there were no

Table 3. Correlations between Q angle, Chippaux-Smirak Index, Arch Index and foot position angle and Fourier Index 56 and Stability Index in eight different positions

	Q angle	CSI	Arch index	FPA
Fall Index	0.461**	0.310**	-0.371**	-0.294*
Eyes open				
Fourier Index 56	0.164	0.061	-0.090	-0.087
Stability Index	0.179	0.051	-0.134	-0.103
Eyes close				
Fourier Index 56	0.460**	0.180	-0.183	-0.131
Stability Index	0.451**	0.171	-0.149	-0.136
Pillow under the feet and eyes open				
Fourier Index 56	0.403**	0.188	-0.124	-0.088
Stability Index	0.439**	0.171	-0.136	-0.100
Pillow under the feet and eyes close				
Fourier Index 56	0.407**	0.115	-0.110	-0.081
Stability Index	0.355*	0.149	-0.096	-0.083
Head turned to the right and eyes close				
Fourier Index 56	0.318*	0.171	-0.143	-0.059
Stability Index	0.342*	0.176	-0.167	-0.078
Head turned to the left and eyes close				
Fourier Index 56	0.350*	0.257*	-0.075	-0.001
Stability Index	0.361*	0.245*	-0.097	-0.023
Head turned to the upside and eyes close				
Fourier Index 56	0.286*	0.176	-0.061	-0.095
Stability Index	0.370**	0.245*	-0.074	-0.105
Head turned to the downside and eyes close				
Fourier Index 56	0.361*	0.252*	-0.199	-0.980
Stability Index	0.374**	0.248*	-0.201	-0.100

SD: Standard deviation; CSI: Chippaux-Smirak Index; FPA: Foot position angle; * *p*<0.05; ** *p*<0.001.

Table 4. Categorical regression analyses with Fall Index as a dependent variable and Q angle, Arch Index, Chippaux-Smirak Index, and foot position angle as independent variables

Variable	Fall index				R ²
	Beta	Standard Error	F	p	
Q angle	0.377	0.106	12.761	0.001**	0.40
Arch Index	-0.324	0.089	13.126	0.001**	
Chippaux-Smirak Index	0.161	0.088	3.380	0.070	
Foot position angle	-0.208	0.090	5.409	0.023*	

* p<0.05; ** p<0.001.

significant differences between the two sexes in terms of the Fall Index, Q angle, CSI, Arch Index, and FPA (p>0.05). Malalignments were as follows: genu varum (n=26), genu valgum (n=6), pes cavus (n=16), pes planus (n=2), and out-toeing (n=10).

In addition, we analyzed the correlation between the Q angle, CSI, Arch Index, and FPA with the Fall Index, eight-position F56, and Stability Index (Table 3). There was a moderate positive correlation between the Fall Index and the Q angle, while a weak positive correlation was found with the CSI and a weak

negative correlation was found with the Arch Index and FPA. Although no correlation of the Q angle with the eyes-open F56 and Stability Index was found, a weak-to-moderate positive correlation was found with these values in the remaining seven positions. There was also a weak positive correlation between the CSI and the F56 and Stability Index in positions where the head turned to the left (HL) and downside (HF). In the position that the head turned to the upside (HB), the CSI was weakly and positively correlated only with the Stability Index. However, no correlation of the Arch

Table 5. Categorical regression analyses with the eyes-closed Fourier Index 56 as a dependent variable and Q angle, Arch Index, Chippaux-Smirak Index, and foot position angle as independent variables. In other regression models with pillow under the feet and eyes-closed Fourier Index 56, head turned to the right and eyes-closed Fourier Index 56, and head turned to the left and eyes-closed Fourier Index 56 as dependent variables and Q angle, Arch Index, and Chippaux-Smirak Index independent variables

Variable	Eyes close - F56				R ²
	Beta	Standard Error	F	p	
Q angle	0.453	0.106	18.185	<.001**	0.260
Arch Index	-0.159	0.098	2.626	0.110	
Chippaux-Smirak Index	0.027	0.102	0.070	0.792	
Foot position angle	-0.079	0.095	0.685	0.411	
Pillow under the feet and eyes close-F56					
Q angle	0.294	0.145	4.096	0.047*	0.107
Arch Index	-0.007	0.101	0.004	0.947	
Chippaux-Smirak Index	0.023	0.112	0.044	0.835	
Foot position angle	-0.047	0.095	0.243	0.623	
Head turned to the right and eyes close-F56					
Q angle	0.297	0.135	4.857	0.031*	0.106
Arch Index	-0.030	0.103	0.088	0.768	
Chippaux-Smirak Index	0.297	0.109	0.067	0.797	
Foot position angle	-0.019	0.096	0.038	0.847	
Head turned to the left and eyes close-F56					
Q angle	0.210	0.139	2.273	0.136	0.138
Arch Index	-0.022	0.118	0.033	0.856	
Chippaux-Smirak Index	0.228	0.104	4.840	0.033*	
Foot position angle	-0.101	0.109	0.853	0.136	

* p<0.05; ** p<0.001.

Index and FPA with the F56 and Stability Index at any positions was found.

Furthermore, the categorical regression models were constructed to evaluate the factors which would affect the Fall Index, F56, and Stability Index. In the first model, the Fall Index was dependent variable, whereas the Q angle, CSI, Arch Index, and FPA were independent variables. The Q angle, Arch Index, and FPA were explained 40% of the variance of the Fall Index ($p=0.001$). The Q angle (Beta: 0.377, $p=0.001$), Arch Index (Beta: -0.324, $p=0.001$), and FPA (Beta: -0.208, $p=0.023$) were independently associated with the Fall Index. These results were demonstrated in Table 4.

In the other regression models, the dependent variables were F56 at eight positions as shown in Table 1 and the Q angle, Arch Index, and FPA were

found to be independent variables. In the models that F56 in eyes-closed, pillow under the feet with eyes-closed positions, and HL were found to be dependent variables, whereas the Q angle, CSI, Arch Index, and FPA were independent variables ($p<0.05$). These results are shown in Table 5. We also found that the Q angle and, to a lesser degree, SCI affected the change in the F56. However, the Arch Index and FPA did not affect F56 significantly.

In final regression models, the Stability Index scores at eight different positions were separately dependent-variable and the Q angle, CSI, Arch Index, and FPA were independent variables. In five of these eight regression models, the p value was <0.05 (Table 6). We found that the Q angle was the most effective variable in the Stability Index, while the effects of CSI and Arch Index were found to be minimal.

Table 6. Categorical regression analyses with the eyes open-Stability Index as dependent variable and Q angle, Arch Index, Chippaux-Smirak Index and foot position angle as independent variables. In other regression models with eyes close-Stability Index pillow under the feet and eyes open-Stability Index, head turned to the upside and eyes close-Stability Index and head turned to the downside and eyes close-Stability Index as dependent variables and Q angle, Arch Index, Chippaux-Smirak Index and as independent variables

Variable	Eyes open-Stability Index				
	Standardized coefficients		F	<i>p</i>	R ²
	Beta	Standard Error			
Q angle	0.180	0.120	2.235	0.139	0.158
Arch Index	-0.295	0.082	12.926	0.001**	
Chippaux-Smirak Index	0.084	0.112	0.562	0.456	
Foot position angle	-0.079	0.093	0.723	0.398	
	Eyes close-Stability Index				
Q angle	0.414	0.104	15.843	<0.001**	0.279
Arch Index	-0.189	0.098	3.694	0.059	
Chippaux-Smirak Index	0.113	0.097	1.362	0.247	
Foot position angle	-0.092	0.090	1.035	0.313	
	Pillow under the feet and eyes open-Stability Index				
Q angle	0.415	0.122	11.468	0.001**	0.223
Arch Index	-0.027	0.107	0.062	0.805	
Chippaux-Smirak Index	0.105	0.111	0.893	0.348	
Foot position angle	-0.071	0.089	0.628	0.431	
	Head turned to the upside and eyes close-Stability Index				
Q angle	0.271	0.134	4.112	0.046*	0.142
Arch Index	-0.056	0.111	0.252	0.617	
Chippaux-Smirak Index	0.167	0.104	2.572	0.113	
Foot position angle	-0.073	0.105	0.488	0.431	
	Head turned to the downside and eyes close-Stability Index				
Q angle	0.231	0.140	2.714	0.104	0.177
Arch Index	-0.082	0.104	0.620	0.434	
Chippaux-Smirak Index	0.264	0.108	5.950	0.017*	
Foot position angle	-0.004	0.104	0.002	0.968	

* $p<0.05$; ** $p<0.001$.

DISCUSSION

Balance is of utmost importance in maintaining postural stability. Balance disorders with impaired motor skills lead to performance loss and an increased risk of falling.^[25] The maintenance of balance depends on several factors including the vestibular system, age, pain, vision, body shape, visual-spatial perception, tactile input, agility, proprioception, and musculoskeletal and neuromuscular systems.^[26] Any defect in this complex may result in imbalance and increase the risk of falling.^[25] It has been shown that the risk of falling increases with increased falling-related risk factors.^[8] In our study, we found a positive correlation between the Fall Index and the Q angle and CSI, and a negative correlation between the Arch Index and FPA. We also showed that the Q angle, Arch Index, and FPA had a statistically significant effect of 40% in the Fall Index. Although the effects of the lower extremity alignment parameters on the risk of falling were not seem to statistically significant, we believe that this effect may be important, considering that falling is multifactorial.

The Q angle is an important indicator of lower extremity alignment disorder.^[26] Numerous studies have shown that an increase in the Q angle causes patellofemoral pain syndrome and, in this syndrome, pain leads to impaired postural control.^[27-29] In a study by Potter et al.,^[30] it was shown that the postural control was impaired in the flexion contracture, while Mahar et al.^[31] reported that the postural control was impaired in the presence of the length difference of the lower limbs. However, the effect of the Q angle on the fall risk in healthy individuals has not been investigated, yet. In our study, we found that the increase in the Q angle also caused an increase in the risk of falling. In addition, we observed that it was the most important factor affecting the risk of falling among the evaluated parameters.

In the joint biomechanics, repetitive proprioceptive inputs constitute the basis of the somatosensory system. The F56 is an indicator of somatosensory disorder; therefore, it provides information about proprioception.^[32] Proprioception plays a key role in the establishment and maintenance of the joint stability. As a result, neuromuscular control is impaired, reflex muscular activities cannot be performed, and the fall risk increases in impaired proprioception.^[33] In our study, we found that, in most of the positions, the Q angle was correlated with the F56 and was the most effective parameter on F56. We considered that one of the reasons of an increased fall risk related to the Q angle, as in our study, may be a proprioceptive disorder

related to its effect on the F56. However, since its impact on the F56 is not very strong, it is not possible to explain this situation with proprioception alone. The Q angle is an indicator of the strength of the quadriceps muscle and an increase in this angle reduces the mechanical effect of the power created by the quadriceps femoris.^[34] It has been shown that this increase may also affect the neuromuscular response and the quadriceps reflex response time.^[34,35] While postural control is achieved, peripheral inputs are united in the central nervous system, and numerous appropriate muscular responses are created. The inability of the increase in the Q angle to generate adequate muscular responses by affecting the quadriceps power may be another reason for its effect on the stability index and, hence, on the risk of falls. In previous studies, quadriceps strengthening exercises have been shown to have a positive effect on balance.^[36,37] Therefore, the fall risk can be reduced by rehabilitation programs directed at the correction of the Q angle.

The foot is the structure contacting directly with the supporting surface and, thus, it plays a major role in all weight-bearing activities and balance control.^[1] The Arch Index, CSI, and FPA are common tools to evaluate the lower extremity alignment. In our study, we found a negative correlation between the Arch Index and FPA with the Fall Index, while there was a positive correlation with the CSI. The reduction in the Arch Index is a sign of pes cavus, indicating that the contact between the foot and the surface is less.^[38] Hertel et al.^[3] reported that the postural control was impaired in cavus feet, as in our study. Maki et al.^[39] showed that, as the foot contact surface decreased, less afferent stimuli were transmitted from the cutaneous receptors. It was also suggested that this reduction in sensory input might lead to the impairment of the maintenance of postural control. However, no relationship was found between the Arch Index and F56, which is an indicator of somatosensory disorders, in our study. This shows that the increase in the risk of falling due to pes cavus is not related to the impairment of the stimulus induced by the cutaneous receptors. In our study, we also found no correlation between the Arch Index and Stability Index, although there was a minimal effect on the Stability Index in only one of eight positions. These findings are similar to the results of Karthikeyan et al.^[1] In pes cavus, pronation becomes restricted due to the hypomobility in the subtalar and midtarsal joints, and the center of gravity of the body shifts medially.^[3,39] As a result, this hypomobility along with the reduced arch angle may be a cause of the increased fall risk.

The CSI is one of the parameters used in the evaluation of the foot medial longitudinal arch (MLA) height and the increase in this index is an indicator of the pronation of the foot.^[3] Cob and Hertel^[3,40] reported that COP excursions increased in the prone feet and, consequently, achieving the stability becomes more difficult. The presence of an association between the foot MLA height and lower extremity injuries has been shown in numerous studies.^[16,17,35] The pronation of the foot is characterized with hypermobility of the midfoot and it is thought that hypermobile joint may lead to neuromuscular problems in maintaining balance.^[41,42] In our study, although the CSI was weakly correlated with the Fall Index, we found no significant effect on the Fall Index and most of the other oscillatory parameters. We believe that this might be due to the small sample size in our study.

The foot progression angle is affected by femoral anteversion and tibiofemoral torsion, and it is an indicator of the rotational malalignment of the lower extremity.^[12,13,43] In previous studies, it was shown that changes in the foot progression angle resulted in various muscular and skeletal problems; however, its effect on the risk of falling has not been investigated, yet.^[44-46] In our study, according to the effects of the FPA on the Fall Index, F56 and Stability Index, we found that it only showed effects on the Fall Index. This suggests that the effect of FPA on the Fall Risk cannot be explained by proprioception. However, further studies are needed to explain the mechanism of this effect.

To the best of our knowledge, there is no study in the literature investigating the effect of lower extremity alignment on the risk of falling in healthy individuals, and this is the first study on this issue. However, further, large-scale studies evaluating other lower extremity alignment factors are required to confirm these findings.

In conclusion, we found that lower extremity malalignment increases the risk of falling. We believe that the risk of falling can be decreased by reducing these malalignments and, thus, fall-related mortality and morbidity can be further reduced. In addition, the correction of the lower extremity malalignment by exercises or orthosis would also reduce the fall risk.

Acknowledgments

We would like to thank Prof. Dr. İmran Kurt Ömürlü for her contribution in the statistical analysis.

Declaration of conflicting interests

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The authors received no financial support for the research and/or authorship of this article.

REFERENCES

1. Karthikeyan G, Jadav Jayraj S, Narayanan V. Effect of forefoot type on postural stability - a cross sectional comparative study. *Int J Sports Phys Ther* 2015;10:213-24.
2. Lihavainen K, Sipilä S, Rantanen T, Sihvonen S, Sulkava R, Hartikainen S. Contribution of musculoskeletal pain to postural balance in community-dwelling people aged 75 years and older. *J Gerontol A Biol Sci Med Sci* 2010;65:990-6.
3. Hertel J, Gay MR, Denegar CR. Differences in Postural Control During Single-Leg Stance Among Healthy Individuals With Different Foot Types. *J Athl Train* 2002;37:129-132.
4. Ambrose AF, Paul G, Hausdorff JM. Risk factors for falls among older adults: a review of the literature. *Maturitas* 2013;75:51-61.
5. Stel VS, Smit JH, Pluijm SM, Lips P. Consequences of falling in older men and women and risk factors for health service use and functional decline. *Age Ageing* 2004;33:58-65.
6. McGuine TA, Greene JJ, Best T, Levenson G. Balance as a predictor of ankle injuries in high school basketball players. *Clin J Sport Med* 2000;10:239-44.
7. Lamb SE, Jørstad-Stein EC, Hauer K, Becker C. Development of a common outcome data set for fall injury prevention trials: the Prevention of Falls Network Europe consensus. *J Am Geriatr Soc* 2005;53:1618-22.
8. Tinetti ME. Clinical practice. Preventing falls in elderly persons. *N Engl J Med*. 2003;348:42-9.
9. Letafatkar A, Zandi S, Khodayi M, Vashmesara JB. Flat Foot Deformity, Q Angle and Knee Pain are Interrelated in Wrestlers. *J Nov Physiother* 2013;3:138.
10. Brattström H. Patella alta in non-dislocating knee joints. *Acta Orthop Scand* 1970;41:578-88.
11. Pezzan P, Sacco I, João S. Foot posture and classification of the plantar arch among adolescent wearers and non-wearers of high-heeled shoes. *Rev Bras Fisioter* 2009;13:398-404.
12. Panjavi B, Mortazavi J. Rotational deformities of the lower limb in children. *Iran J Pediatr* 2007;17:393-39.
13. Cibulka MT, Winters K, Kampwerth T, McAfee B, Payne L, Roeckenhaus T, et al. Predicting foot progression angle during gait using two clinical measures in healthy adults, a preliminary study. *Int J Sports Phys Ther* 2016;11:400-8.
14. Sims EL, Hardaker WM, Queen RM. Gender differences in plantar loading during three soccer-specific tasks. *Br J Sports Med* 2008;42:272-7.
15. Hadadi M, Mazaheri M, Mousavi ME, Maroufi N, Bahramzadeh M, Fardipour S. Effects of soft and semi-rigid ankle orthoses on postural sway in people with and without functional ankle instability. *J Sci Med Sport* 2011;14:370-5.
16. Hreljac A, Marshall RN, Hume PA. Evaluation of lower extremity overuse injury potential in runners. *Med Sci Sports Exerc* 2000;32:1635-41.

17. Bennett JE, Reinking MF, Pluemer B, Pentel A, Seaton M, Killian C. Factors contributing to the development of medial tibial stress syndrome in high school runners. *J Orthop Sports Phys Ther* 2001;31:504-10.
18. Cobb SC, Bazett-Jones DM, Joshi MN, Earl-Boehm JE, James CR. The relationship among foot posture, core and lower extremity muscle function, and postural stability. *J Athl Train* 2014;49:173-80.
19. Tsai LC, Yu B, Mercer VS, Gross MT. Comparison of different structural foot types for measures of standing postural control. *J Orthop Sports Phys Ther* 2006;36:942-53.
20. Hyong IH, Kang JH. Comparison of dynamic balance ability in healthy university students according to foot shape. *J Phys Ther Sci* 2016;28:661-4.
21. Kim HS, Yun DH, Yoo SD, Kim DH, Jeong YS, Yun JS, et al. Balance control and knee osteoarthritis severity. *Ann Rehabil Med* 2011;35:701-9.
22. Greene CC, Edwards TB, Wade MR, Carson EW. Reliability of the quadriceps angle measurement. *Am J Knee Surg* 2001;14:97-103.
23. Aydin E, Turan Y, Tastaban E, Kurt Omurlu I, Sendur OF. Plantar pressure distribution in patients with ankylosing spondylitis. *Clin Biomech (Bristol, Avon)* 2015;30:238-42.
24. Razeghi M, Batt ME. Foot type classification: a critical review of current methods. *Gait Posture* 2002;15:282-91.
25. Citaker S, Kaya D, Yuksel I, Yosmaoglu B, Nyland J, Atay OA, et al. Static balance in patients with patellofemoral pain syndrome. *Sports Health* 2011;3:524-7.
26. Jones K, Barker K. *Human Movement Explained*. Edinburgh, Scotland: Butterworth-Heinemann; 2000.
27. Haim A, Yaniv M, Dekel S, Amir H. Patellofemoral pain syndrome: validity of clinical and radiological features. *Clin Orthop Relat Res* 2006;451:223-8.
28. Loudon JK, Wiesner D, Goist-Foley HL, Asjes C, Loudon KL. Intrarater Reliability of Functional Performance Tests for Subjects With Patellofemoral Pain Syndrome. *J Athl Train* 2002;37:256-261.
29. Aminaka N, Gribble PA. Patellar taping, patellofemoral pain syndrome, lower extremity kinematics, and dynamic postural control. *J Athl Train* 2008;43:21-8.
30. Potter PJ, Kirby RL, MacLeod DA. The effects of simulated knee-flexion contractures on standing balance. *Am J Phys Med Rehabil* 1990;69:144-7.
31. Mahar RK, Kirby RL, MacLeod DA. Simulated leg-length discrepancy: its effect on mean center-of-pressure position and postural sway. *Arch Phys Med Rehabil*. 1985;66:822-4.
32. Lee GE, Bae H, Yoon TS, Kim JS, Yi TI, Park JS. Factors that Influence Quiet Standing Balance of Patients with Incomplete Cervical Spinal Cord Injuries. *Ann Rehabil Med* 2012;36:530-7.
33. Pai YC, Rymer WZ, Chang RW, Sharma L. Effect of age and osteoarthritis on knee proprioception. *Arthritis Rheum* 1997;40:2260-5.
34. Chester R, Smith TO, Sweeting D, Dixon J, Wood S, Song F. The relative timing of VMO and VL in the aetiology of anterior knee pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord* 2008;9:64.
35. de Moura Campos Carvalho-E-Silva AP, Peixoto Leão Almeida G, Oliveira Magalhães M, Renovato França FJ, Vidal Ramos LA, Comachio J, et al. Dynamic postural stability and muscle strength in patellofemoral pain: Is there a correlation? *Knee* 2016;23:616-21.
36. Morag E, Cavanagh PR. Structural and functional predictors of regional peak pressures under the foot during walking. *J Biomech* 1999;32:359-70.
37. Lathinghouse LH, Trimble MH. Effects of isometric quadriceps activation on the Q-angle in women before and after quadriceps exercise. *J Orthop Sports Phys Ther* 2000;30:211-6.
38. Chevidikunnan MF, Al Saif A, Gaowgzeh RA, Mamdouh KA. Effectiveness of core muscle strengthening for improving pain and dynamic balance among female patients with patellofemoral pain syndrome. *J Phys Ther Sci* 2016;28:1518-23.
39. Maki BE, Perry SD, Norrie RG, McIlroy WE. Effect of facilitation of sensation from plantar foot-surface boundaries on postural stabilization in young and older adults. *J Gerontol A Biol Sci Med Sci* 1999;54:281-7.
40. Donatelli RA. *The Biomechanics of the Foot and Ankle*. Philadelphia, PA: F.A. Davis Company; 1996
41. Cobb SC, Tis LL, Johnson BF, Higgie EJ. The effect of forefoot varus on postural stability. *J Orthop Sports Phys Ther* 2004;34:79-85.
42. Era P, Schroll M, Ytting H, Gause-Nilsson I, Heikkinen E, Steen B. Postural balance and its sensory-motor correlates in 75-year-old men and women: a cross-national comparative study. *J Gerontol A Biol Sci Med Sci* 1996;51:53-63.
43. Bryant A, Tinley P, Singer K. Radiographic measurements and planter pressure distribution in normal, hallux valgus and hallux limitus feet. *Foot* 2000;10:18-22.
44. Fuchs R, Staheli LT. Sprinting and intoeing. *J Pediatr Orthop* 1996;16:489-91.
45. Hastings MK, Gelber JR, Isaac EJ, Bohnert KL, Strube MJ, Sinacore DR. Foot progression angle and medial loading in individuals with diabetes mellitus, peripheral neuropathy, and a foot ulcer. *Gait Posture* 2010;32:237-41.
46. Lai YC, Lin HS, Pan HF, Chang WN, Hsu CJ, Renn JH. Impact of foot progression angle on the distribution of plantar pressure in normal children. *Clin Biomech (Bristol, Avon)* 2014;29:196-200.